



EXEMESTANE (Advanced/Metastatic Treatment)

OTHER NAMES: Aromasin®

PURPOSE: Exemestane is a hormonal type medication which may be used for postmenopausal women to treat breast cancer. It works by decreasing the amount of estrogen in the body

ADMINISTRATION: Exemestane is taken by mouth in the form of tablets.
Take once a day at approximately the same time of day.
Tablets take with food or on an empty stomach.

SIDE EFFECTS: Not all patients will experience all side effects.

SIDE EFFECT	MANAGEMENT
<p>Hot flushes: A sudden feeling of warmth which may be accompanied by sweating</p> <p>(common, but usually improves as your body adjusts to the medication)</p>	<ul style="list-style-type: none"> • Dress in layers so you can remove clothing when you get warm. • When possible, lower the room temperature. • Try taking a cool shower or sipping ice water. • If lasting or if troublesome, speak to your doctor or nurse. • Medications may be helpful in managing continuing or bothersome symptoms. Discuss with your doctor.
<p>Muscle, bone or joint pain</p> <p>(common, may occur when you first start taking exemestane)</p>	<ul style="list-style-type: none"> • You may take acetaminophen (Tylenol® and others) or other pain medications such as ASA (aspirin), ibuprofen or others if needed.
<p>Nausea</p> <p>(may occur when you first start taking exemestane-usually mild to moderate)</p>	<ul style="list-style-type: none"> • This rarely requires treatment and may resolve in a short time. • Take exemestane with a meal to lessen stomach upset. • If vomiting occurs and lasts more than 24 hours, contact your doctor or nurse.
<p>Headaches (not common)</p>	<ul style="list-style-type: none"> • You may take acetaminophen (Tylenol® and others) or other pain medications such as ASA (aspirin), ibuprofen or others if needed. • Contact your nurse or doctor if headaches persist.

SIDE EFFECT	MANAGEMENT
Swelling/fluid retention – Hands and feet may swell and you may notice a gain in weight (not common)	<ul style="list-style-type: none"> • Elevate your feet when sitting. • Avoid tight clothing. • Report swelling or weight gain to your doctor.
Other uncommon symptoms you may experience include hair thinning, change in appetite, fatigue, or vaginal dryness	<ul style="list-style-type: none"> • Report these to your doctor.

OTHER IMPORTANT PRECAUTIONS:

- Bone loss or increased blood lipid (cholesterol) levels may occur with exemestane. Discuss with your doctor if necessary.
- Rarely you may experience signs of a blood clot including; shortness of breath, sudden onset of a cough, tenderness or hardness over a vein or calf swelling. Report these symptoms to your doctor immediately.
- Do NOT take any estrogen-containing therapies while taking exemestane.
- Other medications may interact with exemestane. Do not start taking new medications or herbal products without checking with your doctor or pharmacist. Tell your doctor, nurse, or pharmacist if you are taking any other prescription or non-prescription medicine, vitamin, or herbal product.
- Inform other doctors or pharmacists that you are taking exemestane.
- Inform your doctor, nurse, and pharmacist if you develop any new allergies or sensitivities.
- If you experience any symptoms or problems not listed above, inform your doctor, nurse or pharmacist.

STORAGE:

- Store exemestane at room temperature away from heat, direct light, and moisture.
- Whenever possible, return unused medication to the pharmacy from which it was dispensed for proper disposal.

If you have any questions regarding this or any other drug therapy, please phone your cancer centre.

Cross Cancer Institute – (780) 432-8771, Tom Baker Cancer Centre – (403) 521-3764
 Central Alberta Cancer Centre – (403) 343-4802, Grande Prairie Cancer Centre – (780) 538-7288
 Lethbridge Cancer Centre – (403) 329-0633, Medicine Hat Cancer Centre – (403) 529-8817

OR
